

## (12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)


(19) World Intellectual Property Organization  
International Bureau(43) International Publication Date  
10 January 2002 (10.01.2002)

PCT

(10) International Publication Number  
**WO 02/02179 A1**

- (51) International Patent Classification<sup>7</sup>: **A61M 37/00**
- (21) International Application Number: **PCT/US01/20782**
- (22) International Filing Date: **29 June 2001 (29.06.2001)**
- (25) Filing Language: **English**
- (26) Publication Language: **English**
- (30) Priority Data:  
**09/606,909** **29 June 2000 (29.06.2000)** **US**
- (71) Applicant (for all designated States except US): **BECTON, DICKINSON AND COMPANY [US/US]**; Intellectual Property Dept., Mail Code 089, 1 Becton Drive, Franklin Lakes, NJ 07417-1880 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **PETTIS, Ronald, J. [US/US]**; 529 Darby Glen Lane, Durham, NC 27713 (US). **DOWN, James, A. [US/US]**; 814 Oakely Court, Cary, NC 27511 (US). **HARVEY, Noel, G. [US/US]**; 3211 US 70 West, Effland, NC 27243 (US).
- (74) Agents: **SCHMIDT, Richard, D. et al.**; Venable, Baetjer, Howard & Civiletti, LLP, Suite 1000, 1201 New York Avenue, NW, P.O. Box 34385, Washington, DC 20043-9998 (US).
- (81) Designated States (national): **AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.**
- (84) Designated States (regional): **ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).**
- Declaration under Rule 4.17:**  
— of inventorship (Rule 4.17(iv)) for US only

[Continued on next page]

(54) Title: **MICRONEEDLE FOR DELIVERING A SUBSTANCE INTO THE DERMIS**

(57) Abstract: A method for directly delivering whereby a substance is introduced into an intradermal space within mammalian skin which involves administering the substance through at least one small gauge hollow needle having an outlet with an exposed height between 0 and 1 mm. The outlet is inserted into the skin to a depth of between .3 mm and 2 mm such that the delivery of the substance occurs at a depth between .3 mm and 2 mm.

---

**WO 02/02179 A1****Published:**

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

WO 02/02179

PCT/US01/20782

## MICRONEEDLE FOR DELIVERING A SUBSTANCE INTO THE DERMIS

## 5 FIELD OF THE INVENTION

[0001] The present invention relates to methods and devices for administration of substances into the intradermal layer of skin.

## CROSS REFERENCE TO RELATED APPLICATIONS

10 This application is a continuation-in-part of U.S. application No. 09/606,909 filed June 29, 2000.

## BACKGROUND OF THE INVENTION

[0002] The importance of efficiently and safely administering pharmaceutical substances such as diagnostic agents and drugs has long been recognized. Although an important consideration for all pharmaceutical substances, obtaining adequate bioavailability of large molecules such as proteins that have arisen out of the biotechnology industry has recently highlighted this need to obtain efficient and reproducible absorption (Cleland et al., *Curr. Opin. Biotechnol.* 12: 212-219, 2001). The use of conventional needles has long provided one approach for delivering pharmaceutical substances to humans and animals by administration through the skin. Considerable effort has been made to achieve reproducible and efficacious delivery through the skin while improving the ease of injection and reducing patient apprehension and/or pain associated with conventional needles. Furthermore, certain delivery systems eliminate needles entirely, and rely upon chemical mediators or external driving forces such as iontophoretic currents or electroporation or thermal poration or sonophoresis to breach the stratum corneum, the outermost layer of the skin, and deliver substances through the surface of the skin. However, such delivery systems do not reproducibly breach the skin barriers or deliver the pharmaceutical substance to a given depth below the surface of the skin and consequently, clinical results can be variable. Thus, mechanical breach of the stratum corneum such as with needles, is believed to provide the most reproducible method of administration of substances through the surface of the skin, and to provide control and reliability in placement of administered substances.

WO 02/02179

PCT/US01/20782

5

[0003] Approaches for delivering substances beneath the surface of the skin have almost exclusively involved transdermal administration, i.e. delivery of substances through the skin to a site beneath the skin. Transdermal delivery includes subcutaneous, intramuscular or intravenous routes of administration of which, intramuscular (IM) and subcutaneous (SC) injections have been the most commonly used .

10

[0004] Anatomically, the outer surface of the body is made up of two major tissue layers, an outer epidermis and an underlying dermis, which together constitute the skin (for review, see *Physiology, Biochemistry, and Molecular Biology of the Skin, Second Edition*, L.A. Goldsmith, Ed., Oxford University Press, New York, 1991). The epidermis is subdivided into five layers or strata of a total thickness of between 75 and 150  $\mu\text{m}$ . Beneath the epidermis lies the dermis, which contains two layers, an outermost portion referred to as the papillary dermis and a deeper layer referred to as the reticular dermis. The papillary dermis contains vast microcirculatory blood and lymphatic plexuses. In contrast, the reticular dermis is relatively acellular and avascular and made up of dense collagenous and elastic connective tissue. Beneath the epidermis and dermis is the subcutaneous tissue, also referred to as the hypodermis, which is composed of connective tissue and fatty tissue. Muscle tissue lies beneath the subcutaneous tissue.

15

20

[0005] As noted above, both the subcutaneous tissue and muscle tissue have been commonly used as sites for administration of pharmaceutical substances. The dermis, however, has rarely been targeted as a site for administration of substances, and this may be due, at least in part, to the difficulty of precise needle placement into the intradermal space. Furthermore, even though the dermis, in particular, the papillary dermis has been known to have a high degree of vascularity, it has not heretofore been appreciated that one could take advantage of this high degree of vascularity to obtain an improved absorption profile for administered substances compared to subcutaneous administration. This is because small drug molecules are typically rapidly absorbed after administration into the subcutaneous tissue which has been far more easily and predictably targeted than the dermis has been. On the other hand, large molecules such as proteins are typically not well absorbed through the capillary epithelium regardless of the degree of vascularity so that one would not have expected to achieve a significant absorption

30

35

WO 02/02179

PCT/US01/20782

5        advantage over subcutaneous administration by the more difficult to achieve intradermal administration even for large molecules.

10        [0006]     One approach to administration beneath the surface to the skin and into the region of the intradermal space has been routinely used in the Mantoux tuberculin test. In this procedure, a purified protein derivative is injected at a shallow angle to the skin surface using a 27 or 30 gauge needle (Flynn et al, *Chest* 106: 1463-5, 1994). A degree of uncertainty in placement of the injection can, however, result in some false negative test results. Moreover, the test has involved a localized injection to elicit a response at the site of injection and the Mantoux approach has not led to the use of intradermal injection for systemic administration of substances.

20        [0007]     Some groups have reported on systemic administration by what has been characterized as "intradermal" injection. In one such report, a comparison study of subcutaneous and what was described as "intradermal" injection was performed (Autret et al, *Therapie* 46:5-8, 1991). The pharmaceutical substance tested was calcitonin, a protein of a molecular weight of about 3600. Although it was stated that the drug was injected intradermally, the injections used a 4 mm needle pushed up to the base at an angle of 60. This would have resulted in placement of the injectate at a depth of about 3.5 mm and into the lower portion of the reticular dermis or into the subcutaneous tissue rather than into the vascularized papillary dermis. If, in fact, this group injected into the lower portion of the reticular dermis rather than into the subcutaneous tissue, it would be expected that the substance would either be slowly absorbed in the relatively less vascular reticular dermis or diffuse into the subcutaneous region to result in what would be functionally the same as subcutaneous administration and absorption. Such actual or functional subcutaneous administration would explain the reported lack of difference between subcutaneous and what was characterized as intradermal administration, in the times at which maximum plasma concentration was reached, the concentrations at each assay time and the areas under the curves.

35        [0008]     Similarly, Bressolle et al. administered sodium ceftazidime in what was characterized as "intradermal" injection using a 4 mm needle (Bressolle et al. *J. Pharm. Sci.* 82:1175-1178, 1993). This would have resulted in injection to a depth of 4 mm

WO 02/02179

PCT/US01/20782

5 below the skin surface to produce actual or functional subcutaneous injection, although good subcutaneous absorption would have been anticipated in this instance because sodium ceftazidime is hydrophilic and of relatively low molecular weight.

[0009] Another group reported on what was described as an intradermal drug  
10 delivery device (U.S. Patent No. 5,007,501). Injection was indicated to be at a slow rate and the injection site was intended to be in some region below the epidermis, i.e., the interface between the epidermis and the dermis or the interior of the dermis or subcutaneous tissue. This reference, however, provided no teachings that would suggest a selective administration into the dermis nor did the reference suggest any possible  
15 pharmacokinetic advantage that might result from such selective administration.

[0010] Thus there remains a continuing need for efficient and safe methods and devices for administration of pharmaceutical substances.

## 20 SUMMARY OF THE INVENTION

[0011] The present disclosure relates to a new parenteral administration method based on directly targeting the dermal space whereby such method dramatically alters the pharmacokinetics (PK) and pharmacodynamics (PD) parameters of administered substances. By the use of direct intradermal (ID) administration means hereafter  
25 referred to as dermal-access means, for example, using microneedle-based injection and infusion systems (or other means to accurately target the intradermal space), the pharmacokinetics of many substances including drugs and diagnostic substances, which are especially protein and peptide hormones, can be altered when compared to traditional parental administration routes of subcutaneous and intravenous delivery. These findings  
30 are pertinent not only to microdevice-based injection means, but other delivery methods such as needless or needle-free ballistic injection of fluids or powders into the ID space, Mantoux-type ID injection, enhanced iontophoresis through microdevices, and direct deposition of fluid, solids, or other dosing forms into the skin. Disclosed is a method to increase the rate of uptake for parenterally-administered drugs without necessitating IV  
35 access. One significant beneficial effect of this delivery method is providing a shorter  $T_{max}$  (time to achieve maximum blood concentration of the drug). Potential corollary benefits include higher maximum concentrations for a given unit dose ( $C_{max}$ ), higher

WO 02/02179

PCT/US01/20782

5 bioavailability, more rapid uptake rates, more rapid onset of pharmacodynamics or biological effects, and reduced drug depot effects. According to the present invention, improved pharmacokinetics means increased bioavailability, decreased lag time ( $T_{lag}$ ), decreased  $T_{max}$ , more rapid absorption rates, more rapid onset and/or increased  $C_{max}$  for a given amount of compound administered, compared to subcutaneous, intramuscular or  
10 other non-IV parenteral means of drug delivery.

[0012] By bioavailability is meant the total amount of a given dosage that reached the blood compartment. This is generally measured as the area under the curve in a plot of concentration vs. time. By "lag time" is meant the delay between the administration  
15 of a compound and time to measurable or detectable blood or plasma levels.  $T_{max}$  is a value representing the time to achieve maximal blood concentration of the compound, and  $C_{max}$  is the maximum blood concentration reached with a given dose and administration method. The time for onset is a function of  $T_{lag}$ ,  $T_{max}$  and  $C_{max}$ , as all of these parameters influence the time necessary to achieve a blood (or target tissue)  
20 concentration necessary to realize a biological effect.  $T_{max}$  and  $C_{max}$  can be determined by visual inspection of graphical results and can often provide sufficient information to compare two methods of administration of a compound. However, numerical values can be determined more precisely by analysis using kinetic models (as described below) and/or other means known to those of skill in the art.

25 [0013] Directly targeting the dermal space as taught by the invention provides more rapid onset of effects of drugs and diagnostic substances. The inventors have found that substances can be rapidly absorbed and systemically distributed via controlled ID administration that selectively accesses the dermal vascular and lymphatic  
30 microcapillaries, thus the substances may exert their beneficial effects more rapidly than SC administration. This has special significance for drugs requiring rapid onset, such as insulin to decrease blood glucose, pain relief such as for breakthrough cancer pain, or migraine relief, or emergency rescue drugs such as adrenaline or anti-venom. Natural hormones are also released in pulsatile fashion with a rapid onset burst followed by rapid  
35 clearance. Examples include insulin that is released in response to biological stimulus, for example high glucose levels. Another example is female reproductive hormones, which are released at time intervals in pulsatile fashion. Human growth hormone is also

WO 02/02179

PCT/US01/20782

5 released in normal patients in a pulsatile fashion during sleep. This benefit allows better therapy by mimicking the natural body rhythms with synthetic drug compounds. Likewise, it may better facilitate some current therapies such as blood glucose control via insulin delivery. Many current attempts at preparing "closed loop" insulin pumps are hindered by the delay period between administering the insulin and waiting for the  
10 biological effect to occur. This makes it difficult to ascertain in real-time whether sufficient insulin has been given, without overtitrating and risking hypoglycemia. The more rapid PK/PD of ID delivery eliminates much of this type of problem.

[0014] Mammalian skin contains two layers, as discussed above, specifically, the  
15 epidermis and dermis. The epidermis is made up of five layers, the stratum corneum, the stratum lucidum, the stratum granulosum, the stratum spinosum and the stratum germinativum and the dermis is made up of two layers, the upper papillary dermis and the deeper reticular dermis. The thickness of the dermis and epidermis varies from individual to individual, and within an individual, at different locations on the body. For  
20 example, it has been reported that the epidermis varies in thickness from about 40 to about 90  $\mu\text{m}$  and the dermis varies in thickness ranging from just below the epidermis to a depth of from less than 1 mm in some regions of the body to just under 2 to about 4 mm in other regions of the body depending upon the particular study report (Hwang et al., *Ann Plastic Surg* 46:327-331, 2001; Southwood, *Plast. Reconstr. Surg* 15:423-429,  
25 1955; Rushmer et al., *Science* 154:343-348, 1966).

[0015] As used herein, intradermal is intended to mean administration of a substance into the dermis in such a manner that the substance readily reaches the richly vascularized papillary dermis and is rapidly absorbed into the blood capillaries and/or  
30 lymphatic vessels to become systemically bioavailable. Such can result from placement of the substance in the upper region of the dermis, i.e. the papillary dermis or in the upper portion of the relatively less vascular reticular dermis such that the substance readily diffuses into the papillary dermis. It is believed that placement of a substance predominately at a depth of at least about 0.3 mm, more preferably, at least about 0.4  
35 mm and most preferably at least about 0.5 mm up to a depth of no more than about 2.5 mm, more preferably, no more than about 2.0 mm and most preferably no more than about 1.7 mm will result in rapid absorption of macromolecular and/or hydrophobic



WO 02/02179

PCT/US01/20782

5 substances. Placement of the substance predominately at greater depths and/or into the lower portion of the reticular dermis is believed to result in the substance being slowly absorbed in the less vascular reticular dermis or in the subcutaneous region either of which would result in reduced absorption of macromolecular and/or hydrophobic substances. The controlled delivery of a substance in this dermal space below the  
10 papillary dermis in the reticular dermis, but sufficiently above the interface between the dermis and the subcutaneous tissue, should enable an efficient (outward) migration of the substance to the (undisturbed) vascular and lymphatic microcapillary bed (in the papillary dermis), where it can be absorbed into systemic circulation via these microcapillaries without being sequestered in transit by any other cutaneous tissue  
15 compartment.

[0016] Another benefit of the invention is to achieve more rapid systemic distribution and offset of drugs or diagnostic agents. This is also pertinent for many hormones that in the body are secreted in a pulsatile fashion. Many side effects are  
20 associated with having continuous circulating levels of substances administered. A very pertinent example is female reproductive hormones that actually have the opposite effect (cause infertility) when continuously present in the blood. Likewise, continuous and elevated levels of insulin are suspected to down regulate insulin receptors both in quantity and sensitivity.

25 [0017] Another benefit of the invention is to achieve higher bioavailabilities of drugs or diagnostic agents. This effect has been most dramatic for ID administration of high molecular weight substances, especially proteins, peptides, and polysaccharides. The direct benefit is that ID administration with enhanced bioavailability, allows equivalent  
30 biological effects while using less active agent. This results in direct economic benefit to the drug manufacturer and perhaps consumer, especially for expensive protein therapeutics and diagnostics. Likewise, higher bioavailability may allow reduced overall dosing and decrease the patient's side effects associated with higher dosing.

35 [0018] Another benefit of the invention is the attainment of higher maximum concentrations of drugs or diagnostic substances. The inventors have found that substances administered ID are absorbed more rapidly, with bolus administration

WO 02/02179

PCT/US01/20782

5       resulting in higher initial concentrations. This is most beneficial for substances whose efficacy is related to maximal concentration. The more rapid onset allows higher  $C_{Max}$  values to be reached with lesser amounts of the substance. Therefore, the dose can be reduced, providing an economic benefit, as well as a physiological benefit since lesser amounts of the drug or diagnostic agent has to be cleared by the body.

10

[0019]     Another benefit of the invention is no change in systemic elimination rates or intrinsic clearance mechanisms of drugs or diagnostic agents. All studies to date by the applicants have maintained the same systemic elimination rate for the substances tested as via IV or SC dosing routes. This indicates this dosing route has no change in the biological mechanism for systemic clearance. This is an advantageous from a regulatory standpoint, since degradation and clearance pathways need not be reinvestigated prior to filing for FDA approval. This is also beneficial from a pharmacokinetics standpoint, since it allows predictability of dosing regimes. Some substances may be eliminated from the body more rapidly if their clearance mechanism are concentration dependent.

15       Since ID delivery results in higher  $C_{max}$ , clearance rate may be increased, although the intrinsic mechanism remains unchanged.

20

[0020]     Another benefit of the invention is no change in pharmacodynamic mechanism or biological response mechanism. As stated above, administered drugs by the methods taught by the applicants still exert their effects by the same biological pathways that are intrinsic to other delivery means. Any pharmacodynamic changes are related only to the difference patterns of appearance, disappearance, and drug or diagnostic agent concentrations present in the biological system.

25

[0021]     Using the methods of the present invention, pharmaceutical compounds may be administered as a bolus, or by infusion. As used herein, the term "bolus" is intended to mean an amount that is delivered within a time period of less than ten (10) minutes. "Infusion" is intended to mean the delivery of a substance over a time period greater than ten (10) minutes. It is understood that bolus administration or delivery can be carried out with rate controlling means, for example a pump, or have no specific rate controlling means, for example user self-injection.

30

35

WO 02/02179

PCT/US01/20782

5       **[0022]**     Another benefit of the invention is removal of the physical or kinetic barriers  
invoked when drugs passes through and becomes trapped in cutaneous tissue  
compartments prior to systemic absorption. Elimination of such barriers leads to an  
extremely broad applicability to various drug classes. Many drugs administered  
subcutaneously exert this depot effect -- that is, the drug is slowly released from the SC  
10       space, in which it is trapped, as the rate determining step prior to systemic absorption,  
due to affinity for or slow diffusion through the fatty adipose tissue. This depot effect  
results in a lower  $C_{max}$  and longer  $T_{max}$ , compared to ID, and can result in high inter-  
individual variability of absorption. This effect is also pertinent for comparison to  
transdermal delivery methods including passive patch technology, with or without  
15       permeation enhances, iontophoretic technology, sonophoresis, or stratum corneum  
ablation or disruptive methods. Transdermal patch technology relies on drug  
partitioning through the highly impermeable stratum corneum and epidermal barriers.  
Few drugs except highly lipophilic compounds can breach this barrier, and those that do,  
often exhibit extended offset kinetics due to tissue saturation and entrapment of the  
20       drugs. Active transdermal means, while often faster than passive transfer means, are still  
restricted to compound classes that can be moved by charge repulsion or other electronic  
or electrostatic means, or carried passively through the transient pores caused by  
cavitation of the tissue during application of sound waves. The stratum corneum and  
epidermis still provide effective means for inhibiting this transport. Stratum corneum  
25       removal by thermal or laser ablation, abrasive means or otherwise, still lacks a driving  
force to facilitate penetration or uptake of drugs. Direct ID administration by  
mechanical means overcomes the kinetic barrier properties of skin, and is not limited by  
the pharmaceutical or physicochemical properties of the drug or its formulation  
excipients.

30       **[0023]**     Another benefit of the invention is highly controllable dosing regimens. The  
applicants have determined that ID infusion studies have demonstrated dosing profiles  
that are highly controllable and predictable due to the rapid onset and offset kinetics of  
drugs or diagnostic agents delivered by this route. This allows almost absolute control  
35       over the desired dosing regimen when ID delivery is coupled with a fluid control means  
or other control system to regulate metering of the drug or diagnostic agent into the  
body. This single benefit alone is one of the principal goals of most drug or diagnostic

WO 02/02179

PCT/US01/20782

5 agent delivery methods. Bolus ID substance administration as defined previously results  
in kinetics most similar to IV injection and is most desirable for pain relieving  
compounds, mealtime insulin, rescue drugs, erectile dysfunction compounds, or other  
drugs that require rapid onset. Also included would be combinations of substances  
capable of acting alone or synergistically. Extending the ID administration duration via  
10 infusion can effectively mimic SC uptake parameters, but with better predictability.  
This profile is particularly good for substances such as growth hormones, or analgesics.  
Longer duration infusion, typically at lower infusion rates can result in continuous low  
basal levels of drugs that is desired for anticoagulants, basal insulin, and chronic pain  
therapy. These kinetic profiles can be combined in multiple fashion to exhibit almost  
15 any kinetic profile desired. An example would be to pulsatile delivery of fertility  
hormone (LHRH) for pregnancy induction, which requires intermittent peaks every 90  
minutes with total clearance between pulses. Other examples would be rapid peak onset  
of drugs for migraine relief, followed by lower levels for pain prophylaxis.

20 **[0024]** Another benefit of the invention is reduced degradation of drugs and  
diagnostic agents and/or undesirable immunogenic activity. Transdermal methods using  
chemical enhancers or iontophoresis, or sonophoresis or electroporation or thermal  
poration require that a drug pass through the viable epidermal layer, which has high  
metabolic and immunogenic activity. Metabolic conversion of substances in the  
25 epidermis or sequestration by immunoglobulins reduces the amount of drug available for  
absorption. The ID administration circumvents this problem by placing the drug directly  
in the dermis, thus bypassing the epidermis entirely.

30 **[0025]** These and other benefits of the invention are achieved by directly targeting  
absorption by the papillary dermis and by controlled delivery of drugs, diagnostic  
agents, and other substances to the dermal space of skin. The inventors have found that  
by specifically targeting the intradermal space and controlling the rate and pattern of  
delivery, the pharmacokinetics exhibited by specific drugs can be unexpectedly  
improved, and can in many situations be varied with resulting clinical advantage. Such  
35 pharmacokinetics cannot be as readily obtained or controlled by other parenteral  
administration routes, except by IV access.

WO 02/02179

PCT/US01/20782

5        [0026]     The present invention improves the clinical utility of ID delivery of drugs, diagnostic agents, and other substances to humans or animals. The methods employ dermal-access means (for example a small gauge needle, especially microneedles), to directly target the intradermal space and to deliver substances to the intradermal space as a bolus or by infusion. It has been discovered that the placement of the dermal-access  
10        means within the dermis provides for efficacious delivery and pharmacokinetic control of active substances. The dermal-access means is so designed as to prevent leakage of the substance from the skin and improve adsorption within the intradermal space. The pharmacokinetics of hormone drugs delivered according to the methods of the invention have been found to be vastly different to the pharmacokinetics of conventional SC  
15        delivery of the drug, indicating that ID administration according to the methods of the invention will provide improved clinical results. Delivery devices that place the dermal-access means at an appropriate depth in the intradermal space and control the volume and rate of fluid delivery provide accurate delivery of the substance to the desired location without leakage.

20

      [0027]     Disclosed is a method to increase the rate of uptake for parenterally-administered drugs without necessitating IV access. This effect provides a shorter  $T_{max}$ . Potential corollary benefits include higher maximum concentrations for a given unit dose ( $C_{max}$ ), higher bioavailability, more rapid onset of pharmacodynamics or biological  
25        effects, and reduced drug depot effects.

      [0028]     It has also been found that by appropriate depth control of the dermal-access means within the intradermal space that the pharmacokinetics of hormone drugs delivered according to the methods of the invention can, if required, produce similar  
30        clinical results to that of conventional SC delivery of the drug.

      [0029]     The pharmacokinetic profile for individual compounds will vary according to the chemical properties of the compounds. For example, compounds that are relatively large, having a molecular weight of at least 1000 Daltons as well as larger compounds of  
35        at least 2000 Daltons, at least 4000 Daltons, at least 10,000 Daltons and larger and/or hydrophobic compounds are expected to show the most significant changes compared to traditional parenteral methods of administration, such as intramuscular, subcutaneous or

WO 02/02179

PCT/US01/20782

5 subdermal injection. It is expected that small hydrophilic substances, on the whole, will exhibit similar kinetics for ID delivery compared to other methods.

#### DESCRIPTION OF THE DRAWINGS

10 [0030] Figure 1 shows a time course of plasma insulin levels of intradermal versus subcutaneous bolus administration of fast-acting.

[0031] Figure 2 shows a time course of blood glucose levels of intradermal versus subcutaneous bolus administration of fast-acting insulin.

[0032] Figure 3 shows a comparison of bolus ID dosing of fast-acting versus regular insulin.

15 [0033] Figure 4 shows the effects of different intradermal injection depths for bolus dosing of fast-acting insulin on the time course of insulin levels

[0034] Figure 5 shows a comparison of the time course of insulin levels for bolus dosing of long-acting insulin administered subcutaneously or intradermally.

20 [0035] Figure 6 and 7 show a comparison of the pharmacokinetic availability and the pharmacodynamic results of granulocyte colony stimulating factor delivered intradermally with a single needle or three point needle array, subcutaneously, or intravenously.

[0036] Figures 8, 9 and 10 show a comparison of low molecular weight heparin intradermal delivery by bolus, short duration, long duration infusion with comparison to  
25 subcutaneous infusion.

#### DETAILED DESCRIPTION OF THE INVENTION

[0037] The present invention provides a method for therapeutic treatment by delivery of a drug or other substance to a human or animal subject by directly targeting  
30 the intradermal space, where the drug or substance is administered to the intradermal space through one or more dermal-access means incorporated within the device. Substances infused according to the methods of the invention have been found to exhibit pharmacokinetics superior to, and more clinically desirable than that observed for the same substance administered by SC injection.

35 [0038] The dermal-access means used for ID administration according to the invention is not critical as long as it penetrates the skin of a subject to the desired

WO 02/02179

PCT/US01/20782

5       targeted depth within the intradermal space without passing through it. In most cases,  
the device will penetrate the skin and to a depth of about 0.5-2 mm. The dermal-access  
means may comprise conventional injection needles, catheters or microneedles of all  
known types, employed singularly or in multiple needle arrays. The dermal-access  
means may comprise needleless devices including ballistic injection devices. The terms  
10       "needle" and "needles" as used herein are intended to encompass all such needle-like  
structures. The term microneedles as used herein are intended to encompass structures  
smaller than about 30 gauge, typically about 31-50 gauge when such structures are  
cylindrical in nature. Non-cylindrical structures encompass by the term microneedles  
would therefore be of comparable diameter and include pyramidal, rectangular,  
15       octagonal, wedged, and other geometrical shapes. Dermal-access means also include  
ballistic fluid injection devices, powder-jet delivery devices, piezoelectric,  
electromotive, electromagnetic assisted delivery devices, gas-assisted delivery devices,  
of which directly penetrate the skin to provide access for delivery or directly deliver  
substances to the targeted location within the dermal space. By varying the targeted  
20       depth of delivery of substances by the dermal-access means, pharmacokinetic and  
pharmacodynamic (PK/PD) behavior of the drug or substance can be tailored to the  
desired clinical application most appropriate for a particular patient's condition. The  
targeted depth of delivery of substances by the dermal-access means may be controlled  
manually by the practitioner, or with or without the assistance of indicator means to  
25       indicate when the desired depth is reached. Preferably however, the device has  
structural means for controlling skin penetration to the desired depth within the  
intradermal space. This is most typically accomplished by means of a widened area or  
hub associated with the shaft of the dermal-access means that may take the form of a  
backing structure or platform to which the needles are attached. The length of  
30       microneedles as dermal-access means are easily varied during the fabrication process  
and are routinely produced in less than 2 mm length. Microneedles are also a very sharp  
and of a very small gauge, to further reduce pain and other sensation during the injection  
or infusion. They may be used in the invention as individual single-lumen microneedles  
or multiple microneedles may be assembled or fabricated in linear arrays or two-  
35       dimensional arrays as to increase the rate of delivery or the amount of substance  
delivered in a given period of time. Microneedles may be incorporated into a variety of  
devices such as holders and housings that may also serve to limit the depth of

WO 02/02179

PCT/US01/20782

- 5 penetration. The dermal-access means of the invention may also incorporate reservoirs to contain the substance prior to delivery or pumps or other means for delivering the drug or other substance under pressure. Alternatively, the device housing the dermal-access means may be linked externally to such additional components.
- 10 [0039] IV-like pharmacokinetics is accomplished by administering drugs into the dermal compartment in intimate contact with the capillary microvasculature and lymphatic microvasculature. It should be understood that the terms microcapillaries or capillary beds refer to either vascular or lymphatic drainage pathways within the dermal area.
- 15 [0040] While not intending to be bound by any theoretical mechanism of action, it is believed that the rapid absorption observed upon administration into the dermis is achieved as a result of the rich plexuses of blood and lymphatic vessels in the dermis. However, the presence of blood and lymphatic plexuses in the dermis would not by itself
- 20 be expected to produce an enhanced absorption of macromolecules. This is because capillary endothelium is normally of low permeability or impermeable to macromolecules such as proteins, polysaccharides, nucleic acid polymers, substance having polymers attached such as pegylated proteins and the like. Such macromolecules have a molecular weight of at least 1000 Daltons or of a higher molecular weight of at
- 25 least, 2000 Daltons, at least 4000 Daltons, at least 10,000 Daltons or even higher. Furthermore, a relatively slow lymphatic drainage from the interstitium into the vascular compartment would also not be expected to produce a rapid increase in plasma concentration upon placement of a pharmaceutical substance into the dermis.
- 30 [0041] One possible explanation for the unexpected enhanced absorption reported herein is that upon injection of substances so that they readily reach the papillary dermis an increase in blood flow and capillary permeability results. For example, it is known that a pinprick insertion to a depth of 3 mm produces an increase in blood flow and this has been postulated to be independent of pain stimulus and due to tissue release of
- 35 histamine (Arildsson et al., *Microvascular Res.* 59:122-130, 2000). This is consistent with the observation that an acute inflammatory response elicited in response to skin injury produces a transient increase in blood flow and capillary permeability (see



WO 02/02179

PCT/US01/20782

5     *Physiology, Biochemistry, and Molecular Biology of the Skin, Second Edition*, L.A. Goldsmith, Ed., Oxford Univ. Press, New York, 1991, p. 1060; Wilhem, *Rev. Can. Biol.* 30:153-172, 1971). At the same time, the injection into the intradermal layer would be expected to increase interstitial pressure. It is known that increasing interstitial pressure from values (beyond the "normal range") of about -7 to about +2 mmHg distends  
10     lymphatic vessels and increases lymph flow (Skobe et al., *J. Investig. Dermatol. Symp. Proc.* 5:14-19, 2000). Thus, the increased interstitial pressure elicited by injection into the intradermal layer is believed to elicit increased lymph flow and increased absorption of substances injected into the dermis.

15     [0042]     By "improved pharmacokinetics" it is meant that an enhancement of pharmacokinetic profile is achieved as measured, for example, by standard pharmacokinetic parameters such as time to maximal plasma concentration ( $T_{max}$ ), the magnitude of maximal plasma concentration ( $C_{max}$ ) or the time to elicit a minimally detectable blood or plasma concentration ( $T_{lag}$ ). By enhanced absorption profile, it is  
20     meant that absorption is improved or greater as measured by such pharmacokinetic parameters. The measurement of pharmacokinetic parameters and determination of minimally effective concentrations are routinely performed in the art. Values obtained are deemed to be enhanced by comparison with a standard route of administration such as, for example, subcutaneous administration or intramuscular administration. In such  
25     comparisons, it is preferable, although not necessarily essential, that administration into the intradermal layer and administration into the reference site such as subcutaneous administration involve the same dose levels, i.e. the same amount and concentration of drug as well as the same carrier vehicle and the same rate of administration in terms of amount and volume per unit time. Thus, for example, administration of a given  
30     pharmaceutical substance into the dermis at a concentration such as 100 µg/ml and rate of 100 µL per minute over a period of 5 minutes would, preferably, be compared to administration of the same pharmaceutical substance into the subcutaneous space at the same concentration of 100 µg/ml and rate of 100 µL per minute over a period of 5 minutes.

35

[0043]     The enhanced absorption profile is believed to be particularly evident for substances which are not well absorbed when injected subcutaneously such as, for

WO 02/02179

PCT/US01/20782

5 example, macromolecules and/or hydrophobic substances. Macromolecules are, in  
general, not well absorbed subcutaneously and this may be due, not only to their size  
relative to the capillary pore size, it may also be due to their slow diffusion through the  
interstitium because of their size. It is understood that macromolecules can possess  
discrete domains having a hydrophobic and/or hydrophilic nature. In contrast, small  
10 molecules which are hydrophilic are generally well absorbed when administered  
subcutaneously and it is possible that no enhanced absorption profile would be seen  
upon injection into the dermis compared to absorption following subcutaneous  
administration. Reference to hydrophobic substances herein is intended to mean low  
molecular weight substances, for example substances with molecular weights less than  
15 1000 Daltons, which have a water solubility which is low to substantially insoluble

[0044] The above-mentioned PK and PD benefits are best realized by accurate direct  
targeting of the dermal capillary beds. This is accomplished, for example, by using  
microneedle systems of less than about 250 micron outer diameter, and less than 2 mm  
20 exposed length. Such systems can be constructed using known methods of various  
materials including steel, silicon, ceramic, and other metals, plastic, polymers, sugars,  
biological and or biodegradable materials, and/or combinations thereof.

[0045] It has been found that certain features of the intradermal administration  
25 methods provide clinically useful PK/PD and dose accuracy. For example, it has been  
found that placement of the needle outlet within the skin significantly affects PK/PD  
parameters. The outlet of a conventional or standard gauge needle with a bevel has a  
relatively large exposed height (the vertical rise of the outlet). Although the needle tip  
may be placed at the desired depth within the intradermal space, the large exposed height  
30 of the needle outlet causes the delivered substance to be deposited at a much shallower  
depth nearer to the skin surface. As a result, the substance tends to effuse out of the skin  
due to backpressure exerted by the skin itself and to pressure built up from accumulating  
fluid from the injection or infusion. That is, at a greater depth a needle outlet with a  
greater exposed height will still seal efficiently where as an outlet with the same exposed  
35 height will not seal efficiently when placed in a shallower depth within the intradermal  
space. Typically, the exposed height of the needle outlet will be from 0 to about 1 mm.  
A needle outlet with an exposed height of 0 mm has no bevel and is at the tip of the

WO 02/02179

PCT/US01/20782

5        needle. In this case, the depth of the outlet is the same as the depth of penetration of the  
needle. A needle outlet that is either formed by a bevel or by an opening through the  
side of the needle has a measurable exposed height. It is understood that a single needle  
may have more than one opening or outlets suitable for delivery of substances to the  
dermal space.

10

[0046]     It has also been found that by controlling the pressure of injection or infusion  
may avoid the high backpressure exerted during ID administration. By placing a  
constant pressure directly on the liquid interface a more constant delivery rate can be  
achieved, which may optimize absorption and obtain the improved pharmacokinetics.  
15     Delivery rate and volume can also be controlled to prevent the formation of wheals at the  
site of delivery and to prevent backpressure from pushing the dermal-access means out  
of the skin. The appropriate delivery rates and volumes to obtain these effects for a  
selected substance may be determined experimentally using only ordinary skill.  
Increased spacing between multiple needles allows broader fluid distribution and  
20     increased rates of delivery or larger fluid volumes. In addition, it has been found that ID  
infusion or injection often produces higher initial plasma levels of drug than  
conventional SC administration, particularly for drugs that are susceptible to in vivo  
degradation or clearance or for compounds that have an affinity to the SC adipose tissue  
or for macromolecules that diffuse slowly through the SC matrix. This may, in many  
25     cases, allow for smaller doses of the substance to be administered via the ID route.

[0047]     The administration methods useful for carrying out the invention include both  
bolus and infusion delivery of drugs and other substances to humans or animals subjects.  
A bolus dose is a single dose delivered in a single volume unit over a relatively brief  
30     period of time, typically less than about 10 minutes. Infusion administration comprises  
administering a fluid at a selected rate that may be constant or variable, over a relatively  
more extended time period, typically greater than about 10 minutes. To deliver a  
substance the dermal-access means is placed adjacent to the skin of a subject providing  
directly targeted access within the intradermal space and the substance or substances are  
35     delivered or administered into the intradermal space where they can act locally or be  
absorbed by the bloodstream and be distributed systematically. The dermal-access  
means may be connected to a reservoir containing the substance or substances to be

WO 02/02179

PCT/US01/20782

5 delivered. The form of the substance or substances to be delivered or administered  
include solutions thereof in pharmaceutically acceptable diluents or solvents, emulsions,  
suspensions, gels, particulates such as micro- and nanoparticles either suspended or  
dispersed, as well as in-situ forming vehicles of the same. Delivery from the reservoir  
into the intradermal space may occur either passively, without application of the external  
10 pressure or other driving means to the substance or substances to be delivered, and/or  
actively, with the application of pressure or other driving means. Examples of preferred  
pressure generating means include pumps, syringes, elastomer membranes, gas pressure,  
piezoelectric, electromotive, electromagnetic pumping, or Belleville springs or washers  
or combinations thereof. If desired, the rate of delivery of the substance may be variably  
15 controlled by the pressure-generating means. As a result, the substance enters the  
intradermal space and is absorbed in an amount and at a rate sufficient to produce a  
clinically efficacious result.

[0048] As used herein, the term "clinically efficacious result" is meant a clinically  
20 useful biological response including both diagnostically and therapeutically useful  
responses, resulting from administration of a substance or substances. For example,  
diagnostic testing or prevention or treatment of a disease or condition is a clinically  
efficacious result. Such clinically efficacious results include diagnostic results such as  
the measurement of glomerular filtration pressure following injection of inulin, the  
25 diagnosis of adrenocortical function in children following injection of ACTH, the  
causing of the gallbladder to contract and evacuate bile upon injection of  
cholecystokinin and the like as well as therapeutic results, such as clinically adequate  
control of blood sugar levels upon injection of insulin, clinically adequate management  
of hormone deficiency following hormone injection such as parathyroid hormone or  
30 growth hormone, clinically adequate treatment of toxicity upon injection of an antitoxin  
and the like .

[0049] Substances that can be delivered intradermally in accordance with the present  
invention are intended to include pharmaceutically or biologically active substances  
35 including include diagnostic agents, drugs, and other substances which provide  
therapeutic or health benefits such as for example nutraceuticals. Diagnostic substances  
useful with the present invention include macromolecular substances such as, for

WO 02/02179

PCT/US01/20782

5        example, inulin, ACTH (e.g. corticotropin injection), luteinizing hormone-releasing hormone (eg., Gonadorelin Hydrochloride), growth hormone-releasing hormone (e.g. Sermorelin Acetate), cholecystokinin (Sincalide), parathyroid hormone and fragments thereof (e.g. Teriparatide Acetate), thyroid releasing hormone and analogs thereof (e.g. protirelin), secretin and the like.

10

[0050]    Therapeutic substances which can be used with the present invention include Alpha-1 anti-trypsin, Anti-Angiogenesis agents, Antisense, butorphanol, Calcitonin and analogs, Ceredase, COX-II inhibitors, dermatological agents, dihydroergotamine, Dopamine agonists and antagonists, Enkephalins and other opioid peptides, Epidermal growth factors, Erythropoietin and analogs, Follicle stimulating hormone, G-CSF, Glucagon, GM-CSF, granisetron, Growth hormone and analogs (including growth hormone releasing hormone), Growth hormone antagonists, Hirudin and Hirudin analogs such as Hirulog, IgE suppressors, Insulin, insulinotropin and analogs, Insulin-like growth factors, Interferons, Interleukins, Luteinizing hormone, Luteinizing hormone releasing hormone and analogs, Heparins, Low molecular weight heparins and other natural, modified, or syntheic glycoaminoglycans, M-CSF, metoclopramide, Midazolam, Monoclonal antibodies, Peglyated antibodies, Pegylated proteins or any proteins modified with hydrophilic or hydrophobic polymers or additional functional groups, Fusion proteins, Single chain antibody fragments or the same with any combination of attached proteins, macromolecules, or additional functional groups thereof, Narcotic analgesics, nicotine, Non-steroid anti-inflammatory agents, Oligosaccharides, ondansetron, Parathyroid hormone and analogs, Parathyroid hormone antagonists, Prostaglandin antagonists, Prostaglandins, Recombinant soluble receptors, scopolamine, Serotonin agonists and antagonists, Sildenafil, Terbutaline, Thrombolytics, Tissue plasminogen activators, TNF - , and TNF - antagonist, the vaccines, with or without carriers/adjuvants, including prophylactics and therapeutic antigens (including but not limited to subunit protein, peptide and polysaccharide, polysaccharide conjugates, toxoids, genetic based vaccines, live attenuated, reassortant, inactivated, whole cells, viral and bacterial vectors) in connection with, addiction, arthritis, cholera, cocaine addiction, diphtheria, tetanus, HIB, Lyme disease, meningococcus, measles, mumps, rubella, varicella, yellow fever, Respiratory syncytial virus, tick borne japanese encephalitis, pneumococcus, streptococcus, typhoid, influenza, hepatitis, including

35

WO 02/02179

PCT/US01/20782

5 hepatitis A, B, C and E, otitis media, rabies, polio, HIV, parainfluenza, rotavirus, Epstein  
Barr Virus, CMV, chlamydia, non-typeable haemophilus, moraxella catarrhalis, human  
papilloma virus, tuberculosis including BCG, gonorrhoea, asthma, atherosclerosis  
10 malaria, E-coli, Alzheimer's Disease, H. Pylori, salmonella, diabetes, cancer, herpes  
simplex, human papilloma and the like other substances including all of the major  
therapeutics such as agents for the common cold, Anti-addiction, anti-allergy, anti-  
emetics, anti-obesity, antiosteoporotic, anti-infectives, analgesics, anesthetics,  
anorexics, antiarthritics, antiasthmatic agents, anticonvulsants, anti-depressants,  
antidiabetic agents, antihistamines, anti-inflammatory agents, antimigraine preparations,  
15 antinausea preparations, antinauseants, antineoplastics, antiparkinsonism drugs,  
antipruritics, antipsychotics, antipyretics, anticholinergics, benzodiazepine antagonists,  
vasodilators, including general, coronary, peripheral and cerebral, bone stimulating  
agents, central nervous system stimulants, hormones, hypnotics, immunosuppressives,  
muscle relaxants, parasympatholytics, parasympathomimetics, prostaglandins, proteins,  
20 peptides, polypeptides and other macromolecules, psychostimulants, sedatives, and  
sexual hypofunction and tranquilizers.

[0051] Pharmacokinetic analysis of insulin infusion data was carried out as follows.  
Stepwise nonlinear least-squares regression was used to analyze the insulin  
concentration-time data from each individual animal. Initially, an empirical  
25 biexponential equation was fit to the insulin concentration-time data for the negative  
control condition. This analysis assumed first-order release of residual insulin, and  
recovered parameters for the first-order rate constant for release, the residual insulin  
concentration at the release site, a lag time for release, and a first-order rate constant for  
elimination of insulin from the systemic circulation. The parameters recovered in this  
30 phase of the analysis are of no intrinsic importance, but merely account for the fraction  
of circulating insulin derived from endogenous sources.

[0052] The second step of the analysis involved fitting an explicit compartmental  
model to the insulin concentration-time data during and after subcutaneous or  
35 intradermal infusion. The scheme upon which the mathematical model was based is  
shown in the upper part of Figure 1.[PK/PD model fig]. Infusion of insulin proceeded  
from  $t = 0$  to  $t = 240$  min; after a lag time ( $t_{lag,2}$ ), absorption from the infusion site was

WO 02/02179

PCT/US01/20782

5 mediated by a first-order process governed by the absorption rate constant  $k_a$ . Insulin absorbed into the systemic circulation distributed into an apparent volume  $V$  contaminated by an unknown fractional bioavailability  $F$ , and was eliminated according to a first-order rate constant  $K$ . The fitting routine recovered estimates of  $t_{lag,2}$ ,  $k_a$ ,  $V/F$ , and  $K$ ; parameters associated with the disposition of endogenous insulin ( $C_R$ ,  $t_{lag,1}$ ,  $k_R$ ),  
 10 which were recovered in the first step of the analysis, were treated as constants.

[0053] Parameter estimates are reported as mean  $\pm$  SD. The significance of differences in specific parameters between the two different modes of insulin administration (subcutaneous versus intradermal infusion) was assessed with the paired  
 15 Student's  $t$ -test.

Pharmacodynamic analysis of insulin infusion data was calculated as follows. Plasma concentrations of glucose were used as a surrogate for the pharmacological effect of insulin. The change in response variable  $R$  (plasma glucose concentration) with respect  
 20 to time  $t$  was modeled as

$$\frac{dR}{dt} = k_{in} - E \cdot k_{out}$$

[0054] where  $k_{in}$  is the zero-order infusion of glucose,  $k_{out}$  is the first-order rate constant mediating glucose elimination, and  $E$  is the effect of insulin according to the  
 25 sigmoid Hill relationship

$$E = \frac{E_{max} \cdot C^\gamma}{EC_{50}^\gamma + C^\gamma}$$

[0055] in which  $M_{ax}$  is the maximal stimulation of  $k_{out}$  by insulin,  $EC_{50}$  is the insulin concentration at which stimulation of  $k_{out}$  is half maximal,  $C$  is the concentration of  
 30 insulin, and  $\gamma$  is the Hill coefficient of the relationship. Initial modeling efforts utilized the plasma concentration of insulin as the mediator of pharmacological response. However, this approach did not capture the delay in response of plasma glucose to increasing concentrations of plasma insulin. Therefore, an effect-compartment modeling  
 35 approach was finally adopted in which the effect of insulin was mediated from a

WO 02/02179

PCT/US01/20782

5       hypothetical effect compartment peripheral to the systemic pharmacokinetic compartment

10       [0056]     The pharmacodynamic analysis was conducted in two steps. In the first step of the analysis, initial estimates of the pharmacokinetic parameters associated with the disposition of glucose ( $C_{out}$  and the volume of distribution of glucose,  $V_{glucose}$ ) were determined from the glucose concentration-time data in the negative control condition. The full integrated pharmacokinetic-pharmacodynamic model then was fit simultaneously to the glucose concentration-time data from the negative control condition and each insulin delivery condition for each animal (i.e., two sets of  
15       pharmacodynamic parameters were obtained for each animal: one from the simultaneous analysis of the subcutaneous insulin infusion/negative control data, and one from the simultaneous analysis of the intradermal insulin infusion/negative control data). In all pharmacodynamic analyses, the parameters governing insulin disposition obtained during pharmacokinetic analysis of insulin concentration-time data from each animal  
20       were held constant.

[0057]     All other pharmacokinetic analyses were calculated using non-compartmental methods using similar software programs and techniques known in the art..

25       [0058]     Having described the invention in general, the following specific but not limiting examples and reference to the accompanying Figures set forth various examples for practicing the dermal accessing, direct targeting drug administration method and examples of dermal administered drug substances providing improved PK and PD effects.

30       [0059]     A representative example of dermal-access microdevice comprising a single needle were prepared from 34 gauge steel stock (MicroGroup, Inc., Medway, MA) and a single 28° bevel was ground using an 800 grit carborundum grinding wheel. Needles were cleaned by sequential sonication in acetone and distilled water, and flow-checked  
35       with distilled water. Microneedles were secured into small gauge catheter tubing (Maersk Medical) using UV-cured epoxy resin. Needle length was set using a mechanical indexing plate, with the hub of the catheter tubing acting as a depth-limiting



WO 02/02179

PCT/US01/20782

5 control and was confirmed by optical microscopy. For experiments using needles of various lengths, the exposed needle lengths were adjusted to 0.5, 0.8, 1, 2 or 3 mm using the indexing plate. Connection to the fluid metering device, either pump or syringe, was via an integral Luer adapter at the catheter inlet. During injection, needles were inserted perpendicular to the skin surface, and were either held in place by gentle hand pressure  
10 for bolus delivery or held upright by medical adhesive tape for longer infusions. Devices were checked for function and fluid flow both immediately prior to and post injection. This Luer Lok single needle catheter design is hereafter designated SS1\_34.

[0060] Yet another dermal-access array microdevices was prepared consisting of 1”  
15 diameter disks machined from acrylic polymer, with a low volume fluid path branching to each individual needle from a central inlet. Fluid input was via a low volume catheter line connected to a Hamilton microsyringe, and delivery rate was controlled via a syringe pump. Needles were arranged in the disk with a circular pattern of 15 mm diameter. Three-needle and six-needle arrays were constructed, with 12 and 7 mm  
20 needle-to-needle spacing, respectively. All array designs used single-bevel, 34 G stainless steel microneedles of 1 mm length. The 3-needle 12mm spacing catheter-design is hereafter designated SS3\_34B, 6-needle 7mm spacing catheter-design is hereafter designated SS6\_34A.

25 [0061] Yet another dermal-access array microdevices was prepared consisting of 11mm diameter disks machined from acrylic polymer, with a low volume fluid path branching to each individual needle from a central inlet. Fluid input was via a low volume catheter line connected to a Hamilton microsyringe, and delivery rate was controlled via a syringe pump. Needles were arranged in the disk with a circular pattern  
30 of about 5 mm diameter. Three-needle arrays of about 4 mm spacing connected to a catheter as described above. These designs are hereafter designated SS3S\_34\_1, SS3C\_34\_2, and SS3S\_34\_3 for 1mm, 2mm, and 3mm needle lengths respectively.

[0062] Yet another dermal-access ID infusion device was constructed using a  
35 stainless steel 30 gauge needle bent at near the tip at a 90-degree angle such that the available length for skin penetration was 1-2 mm. The needle outlet (the tip of the needle) was at a depth of 1.7-2.0 mm in the skin when the needle was inserted and the

WO 02/02179

PCT/US01/20782

5 total exposed height of the needle outlet 1.0-1.2 mm This design is hereafter designated SSB1\_30.

[0063] EXAMPLE I

[0064] Slow-infusion ID insulin delivery was demonstrated in swine using a hollow,  
10 silicon-based single-lumen microneedle (2 mm total length and 200 X 100  $\mu$ m OD, corresponding to about 33 gauge) with an outlet 1.0  $\mu$ m from the tip (100  $\mu$ m exposed height), was fabricated using processes known in the art (US Patent No. 5,928,207) and mated to a microbore catheter (Disetronic). The distal end of the microneedle was placed into the plastic catheter and cemented in place with epoxy resin to form a depth-limiting hub. The needle outlet was positioned approximately 1 mm beyond the epoxy  
15 hub, thus limiting penetration of the needle outlet into the skin to approximately 1 mm., which corresponds to the depth of the intradermal space in swine.. The catheter was attached to a MiniMed 507 insulin pump for control of fluid delivery. The distal end of the microneedle was placed into the plastic catheter and cemented in place with epoxy  
20 resin to form a depth-limiting hub. The needle outlet was positioned approximately 1 mm beyond the epoxy hub, thus limiting penetration of the needle outlet into the skin to approximately 1 mm., which corresponds to the depth of the intradermal space in swine. The patency of the fluid flow path was confirmed by visual observation, and no obstructions were observed at pressures generated by a standard 1-cc syringe. The  
25 catheter was connected to an external insulin infusion pump (MiniMed 507) via the integral Luer connection at the catheter outlet. The pump was filled with Humalog™ (Lispro) insulin (Eli Lilly, Indianapolis, IN) and the catheter and microneedle were primed with insulin according to the manufacturer's instructions. Sandostatin® (Sandoz, East Hanover, NJ) solution was administered via IV infusion to anesthetized swine to  
30 suppress basal pancreatic function and insulin secretion. After a suitable induction period and baseline sampling, the primed microneedle was inserted perpendicular to the skin surface in the flank of the animal up to the hub stop. Insulin infusion at a rate of 2 U/hr was used and maintained for 4 hr. Blood samples were periodically withdrawn and analyzed for serum insulin concentration and blood glucose values. Baseline insulin  
35 levels before infusion were at the background detection level of the assay. After initiation of the infusion, serum insulin levels showed an increase that was

WO 02/02179

PCT/US01/20782

5 commensurate with the programmed infusion rates. Blood glucose levels also showed a  
corresponding drop relative to negative controls (NC) without insulin infusion and this  
drop was improved relative to conventional SC infusion. In this experiment, the  
microneedle was demonstrated to adequately breach the skin barrier and deliver a drug  
*in vivo* at pharmaceutically relevant rates. The ID infusion of insulin was demonstrated  
10 to be a pharmacokinetically acceptable administration route, and the pharmacodynamic  
response of blood glucose reduction was also demonstrated. Calculated PK parameters  
for ID infusion indicate that insulin is absorbed much faster than via than SC  
administration. Absorption from the ID space begins almost immediately: the lag time  
prior to absorption ( $t_{lag}$ ) was 0.88 vs. 13.6 min for ID and SC respectively. Also the rate  
15 of uptake from the administration site is increased by approximately 3-fold,  $k_a = 0.0666$   
vs.  $0.0225 \text{ min}^{-1}$  for ID and SC respectively. The bioavailability of insulin delivered by  
ID administration is increased approximately 1.3 fold greater than SC administration.

[0065] EXAMPLE II Bolus delivery of Lilly Lispro fast acting insulin was  
20 performed using ID and SC bolus administration. The ID injection microdevice was  
dermal access array design SS3\_34. 10 international insulin units (U) corresponding to  
100 uL volume respectively, were administered to diabetic Yucatan Mini swine. Test  
animals had been previously been rendered diabetic by chemical ablation of pancreatic  
islet cells, and were no longer able to secrete insulin. Test animals received their insulin  
25 injection either via the microneedle array or via a standard 30 G X ½ in. needle inserted  
laterally into the SC tissue space. Circulating serum insulin levels were detected using a  
commercial chemiluminescent assay kit (Immulite, Los Angeles, CA) and blood  
glucose values were determined using blood glucose strips. ID injections were  
accomplished via hand pressure using an analytical microsyringe and were administered  
30 over approximately 60 sec. By comparison, SC dosing required only 2-3 sec. Referring  
to Figure 1, it is shown that serum insulin levels after bolus administration demonstrate  
more rapid uptake and distribution of the injected insulin when administered via the ID  
route. The time to maximum concentration ( $T_{max}$ ) is shorter and the maximum  
concentration obtained ( $C_{max}$ ) is higher for ID vs. SC administration. In addition, Figure  
35 2 also demonstrates the pharmacodynamic biological response to the administered  
insulin, as measured by the decrease in blood glucose (BG), showed faster and greater  
changes in BG since more insulin was available early after ID administration.

WO 02/02179

PCT/US01/20782

5

**[0066] EXAMPLE III**

**[0067]** Lilly Lispro is regarded as fact acting insulin, and has a slightly altered protein structure relative to native human insulin.. Hoechst regular insulin, maintains the native human insulin protein structure that is chemically similar, but has slower uptake than Lispro when administered by the traditional SC route. Both insulin types were administered in bolus via the ID route to determine if any differences in uptake would be discernable by this route. 5U of either insulin type were administered to the ID space using dermal access microdevice design SS3\_34. The insulin concentration verses time data shown in Figure 3. When administered by the ID route the PK profiles for regular and fast-acting insulin were essentially identical, and both insulin types exhibited faster uptake than Lispro given by the traditional SC route. This is evidence that the uptake mechanism for ID administration is less affected by minor biochemical changes in the administered substance, and that ID delivery provides an advantageous PK uptake profile for regular insulin that is superior to SC administered fast-acting insulin.

20

**[0068] EXAMPLE IV**

**[0069]** Bolus delivery of Lilly Lispro fast-acting insulin via microneedle arrays having needles of various lengths was conducted to demonstrate that the precise deposition of drug into the dermal space is necessary to obtain the PK advantages and distinctions relative to SC. Thus, 5U of Lilly Lispro fast-acting insulin was administered using dermal access design SS3\_34. Additional microdevices of the same needle array configuration were fabricated whereby exposed needle lengths of the microdevice array were lengthened to include arrays with needles lengths of 2 and 3 mm. The average total dermal thickness in Yucatan Mini swine ranges from 1.5-2.5 mm. Therefore insulin deposition is expected to be into the dermis, approximately at the dermal/SC interface, and below the dermis and within the SC for 1mm, 2mm, and 3mm length needles respectively. Bolus insulin administration was as described in EXAMPLE II.. Average insulin concentrations verses time are shown in Figure 4. The data clearly shows as microneedle length is increased, the resulting PK profile begins to more closely resemble SC administration. This data demonstrates the benefits of directly targeting the dermal space, such benefits include rapid uptake and distribution, and high initial concentrations. Since the data are averages of multiple examples, they do not show the

35

WO 02/02179

PCT/US01/20782

5 increased inter-individual variability in PK profiles from longer 2 and 3mm  
microneedles. This data demonstrates that since skin thickness may vary both between  
individuals and even within a single individual, shorter needle lengths that accurately  
target the dermal space are more reproducible in their PK profile since they are  
depositing the drug more consistently in the same tissue compartment. This data  
10 demonstrates longer microneedles that deposit or administer substances deeper into the  
dermal space, or partially or wholly into the SC space, mitigate or eliminate the PK  
advantages in comparison to shallow, directly targeted administrations to the highly  
vascularized dermal region.

15 [0070] EXAMPLE V

[0071] Bolus delivery of Lantus long-acting insulin was delivered via the ID route .  
Lantus is an insulin solution that forms microprecipitates at the administration site upon  
injection. These microparticulates undergo slow dissolution within the body to provide  
20 (according to the manufacturer's literature) a more stable low level of circulating insulin  
than other current long-acting insulin such as crystalline zinc precipitates (e.g. Lente,  
NPH). Lantus insulin (10 U dose, 100 uL) was administered to diabetic Yucatan Mini  
pigs using the dermal access design SS3\_34 and by the standard SC method as  
previously described. Referring to Figure 5, when administered via the ID route, similar  
PK profiles were obtained relative to SC. Minor distinctions include a slightly higher  
25 "burst" immediately after the ID insulin delivery. This demonstrates that the uptake of  
even very high molecular weight compounds or small particles is achievable via ID  
administration. More importantly this supports the fact that the biological clearance  
mechanism in the body is not appreciably changed by the administration route, nor is the  
way in which that the drug substance is utilized. This is extremely important for drugs  
30 compounds that have a long circulating half-life (examples would be large soluble  
receptor compounds or other antibodies for cancer treatment, or chemically modified  
species such as PEGylated drugs).

[0072] EXAMPLE VI

35 [0073] Bolus ID delivery of human granulocyte colony stimulating factor (GCSF)  
(Neupogen) was administered via dermal access microdevice designs SS3\_34B (array)  
or SS1\_34 (single needle) to Yucatan minipigs. Delivery rate was controlled via a

WO 02/02179

PCT/US01/20782

5 Harvard syringe pump and was administered over a 1-2.5 min period. Figure 6 shows the PK availability of GCSF in blood plasma as detected by an ELISA immunoassay specific for GCSF. Administration via IV and SC delivery was performed as controls. Referring to Figure 6 bolus ID delivery of GCSF shows the more rapid uptake associated with ID delivery.  $C_{max}$  is achieved at approximately 30-90 minutes vs. 120 min for SC.

10 Also the bioavailability is dramatically increased by an approximate factor of 2 as evidenced by the much higher area under the curve (AUC). Circulating levels of GCSF are detectable for an extended period, indicating that ID delivery does not alter the intrinsic biological clearance mechanism or rate for the drug. These data also show that device design has minimal effect on the rapid uptake of drug from the ID space. The data

15 referred to in Figure 7 also shows the degree and time course of white blood cell expansion as a result of GCSF administration with respect to a negative control (no GCSF administered). White blood cell (WBC) counts were determined by standard cytometric clinical veterinary methods ID delivery exhibits the same clinically significant biological outcomes. Although all delivery means give approximately equal

20 PD outcomes, this data suggests ID delivery could enable use half the dose to achieve essentially the same physiological result in comparison to SC, due to approximately 2-fold bioavailability increase.

#### [0074] EXAMPLE VII

25 An ID administration experiment was conducted using a peptide drug entity: human parathyroid hormone 1-34 (PTH). PTH was infused for a 4 h period, followed by a 2 h clearance. Control SC infusion was through a standard 31-gauge needle inserted into the SC space lateral to the skin using a "pinch-up" technique. ID infusion was through dermal access microdevice design SSB1\_30 (a stainless steel 30-gauge needle bent at the

30 tip at a 90° angle such that the available length for skin penetration was 1-2 mm). The needle outlet (the tip of the needle) was at a depth of 1.7-2.0 mm in the skin when the needle was inserted. A 0.64 mg/mL PTH solution was infused at a rate of 75  $\mu$ L/hr. Flow rate was controlled via a Harvard syringe pump. Weight normalized PTH plasma levels are shown in Figure XX. {The weight normalized delivery profiles show a larger

35 area under the curve (AUC) indicating higher bioavailability, higher peak values at earlier sampling timepoints (e.g. 15 and 30 min) indicating more rapid onset from ID

WO 02/02179

PCT/US01/20782

5 delivery, and rapid decrease following termination of infusion (also indicative of rapid uptake without a depot effect).}

10 [0075] The above examples and results demonstrate the inventive delivery method using multi-point array ID administration and single needle ID administration results in more rapid uptake with higher  $C_{\max}$  than SC injection. ID uptake and distribution is ostensibly unaffected by device geometry parameters, using needle lengths of about 0.5 to about 1.7mm, needle number and needle spacing. No concentration limit for biological absorption was found and PK profiles were dictated principally by the concentration-based delivery rate. The primary limitations of ID administration are the total volume and volumetric infusion-rate limits for leak-free instillation of exogenous substances into a dense tissue compartment. Since absorption of drugs from the ID space appears to be insensitive to both device design and volumetric infusion rate, numerous formulation/device combinations can be used to overcome this limitations and provide the required or desired therapeutic profiles. For example, volume limited dosing regimens can be circumvented either by using more concentrated formulations or increasing the total number of instillation sites. In addition, effective PK control is obtained by manipulating infusion or administration rate of substances.

25 [0076] In general, ID delivery as taught by the methods described hereto via dermal access microneedle devices provides a readily accessible and reproducible parenteral delivery route, with high bioavailability, as well as the ability to modulate plasma profiles by adjusting the device infusion parameters, since uptake is not rate-limited by biological uptake parameters.

30 [0077] In the previously described examples, the methods practiced by the invention demonstrate the ability to deliver a drug in vivo with greatly improved pharmaceutically relevant rates. This data indicates an improved pharmacological result for ID administration as taught by the methods described of other drugs in humans would be expected according to the methods of the invention.

35

WO 02/02179

PCT/US01/20782

5       **[0078]**     All references cited in this specification are hereby incorporated by reference.  
The discussion of the references herein is intended merely to summarize the assertions  
made by their authors and no admission is made that any reference constitutes prior art  
relevant to patentability. Applicants reserve the right to challenge the accuracy and  
pertinency of the cited references.

10



WO 02/02179

PCT/US01/20782

WHAT IS CLAIMED IS:

- 5           1. A method for directly delivering a substance into an intradermal space within mammalian skin comprising administering the substance through at least one small gauge hollow needle having an outlet with an exposed height between 0 and 1 mm, said outlet being inserted into the skin to a depth of between .3 mm and 2 mm, such that delivery of the substance occurs at a depth between .3 mm and 2 mm.
- 10           2. The method according to claim 1 wherein the delivered substance has improved pharmacokinetics compared to pharmacokinetics after subcutaneous injection.
- 15           3. The method of claim 1 wherein the administration is through at least one small gauge hollow needle.
- 20           4. The method of claim 1 wherein the needle has an outlet with an exposed height between 0 and 1 mm.
- 25           5. The method of Claim 1 wherein injecting comprises inserting the needle to a depth which delivers the substance at least about 0.3 mm below the surface to no more than about 2 mm below the surface.
- 30           6. The method of Claim 1 wherein administering comprises inserting the needle into the skin to a depth of at least about 0.3 mm and no more than about 2 mm.
- 35           7. The method of claim 2 wherein the improved pharmacokinetics is increased bioavailability of the substance.
8. The method of claim 2 wherein the improved pharmacokinetics is a decrease in  $T_{max}$ .
9. The method of claim 2 wherein the improved pharmacokinetics is an increase in  $C_{max}$ .

WO 02/02179

PCT/US01/20782

5           10. The method of claim 2 wherein the improved pharmacokinetics is a decrease in  
T<sub>lag</sub>.

11. The method of claim 2 wherein the improved pharmacokinetics is enhanced  
absorption rate.

10

12. The method of claim 1 wherein the substance is administered over a time period  
of not more than ten minutes.

13. The method of claim 1 wherein the substance is administered over a time period  
15 of greater than ten minutes.

14. The method of claim 1 wherein the substance is a peptide or protein.

15. The method of claim 1 wherein the substance is administered at a rate between 1  
20 nL/min. and 200 mL/ min.

16. The method of claim 1 wherein said substance is a hormone.

17. The method of claim 14 wherein said protein or peptide is selected from the  
25 group consisting of insulin, granulocyte stimulating factor and PTH.

18. The method of claim 1 wherein said substance is a nucleic acid.

19. The method of claim 1 wherein the substance has a molecular weight of less than  
30 1000 daltons.

20. The method of claim 1 wherein the substance has a molecular weight greater than  
1000 daltons.

35           21. The method of claim 1 wherein said substance is hydrophobic.

22. The method of claim 1 wherein said substance is hydrophilic.

WO 02/02179

PCT/US01/20782

5           23. The method of claim 1 wherein the needle(s) are inserted substantially  
perpendicularly to the skin.

          24. A method of administering a pharmaceutical substance comprising injecting or  
infusing the substance intradermally through one or more microneedles having a length and  
10       outlet suitable for selectively delivering the substance into the dermis to obtain absorption of  
the substance in the dermis.

          25. The method of Claim 24 wherein absorption of the substance in the dermis  
produces improved systemic pharmacokinetics compared to subcutaneous administration.

15

          26. The method of Claim 25 wherein the improved pharmacokinetics is increased  
bioavailability.

          27. The method of Claim 25 wherein the improved pharmacokinetics is decreased  
20        $T_{max}$ .

          28. The method of claim 25 wherein the improved pharmacokinetics is an increase in  
 $C_{max}$ .

25       29. The method of claim 25 wherein the improved pharmacokinetics is a decrease in  
 $T_{lag}$ .

          30. The method of claim 25 wherein the improved pharmacokinetics is an enhanced  
absorption rate.

30

          31. The method of claim 24 wherein the length of the microneedle is from about 0.5  
mm to about 1.7 mm.

          32. The method of Claim 24 wherein the microneedle is a 30 to 34 gauge needle

35

          33. The method of Claim 24 wherein the microneedle has an outlet of from 0 to 1  
mm.

WO 02/02179

PCT/US01/20782

- 5           34. The method of Claim 24 wherein the microneedle is configured in a delivery device which positions the microneedle perpendicular to skin surface.
35. The method of Claim 24 wherein the microneedle needle is contained in an array of microneedles needles.
- 10           36. The method of Claim 35 wherein the array comprises 3 microneedles.
37. The method of Claim 35 wherein the array comprises 6 microneedles.
- 15           38. A microneedle for intradermal injection of a pharmaceutical substance, wherein the microneedle has a length and outlet selected for its suitability for specifically delivering the substance into the dermis.
39. The microneedle according to claim 38 wherein the length of the microneedle is
- 20           from about 0.5 mm to about 1.7 mm.
40. The microneedle of Claim 38 which is a 30 to 34 gauge needle
41. The microneedle of Claim 38 which has an outlet of from 0 to 1 mm
- 25           42. The microneedle of Claim 38 which is configured in a delivery device which positions the microneedle perpendicular to skin surface.
43. The microneedle of Claim 42 which is in an array of microneedles needles.
- 30           44. The microneedle of Claim 43 wherein the array comprises 3 microneedles.
45. The microneedle of Claim 43 wherein the array comprises 6 microneedles.

WO 02/02179

PCT/US01/20782


- 5           46. A method for delivering a bioactive substance to a subject comprising :  
contacting the skin of the subject with a device having a dermal-access means for accurately  
targeting the dermal space of the subject     with an efficacious amount of the bioactive  
substance.
- 10           47. The method of claim 46 wherein the pharmacokinetics of the bioactive substance  
is improved relative to the pharmacokinetics of the substance when administered  
subcutaneously.
48. The method of claim 47 wherein the improved pharmacokinetics is an increase in  
15   bioavailability.
49. The method of claim 47 wherein the improved pharmacokinetics is a decrease in  
 $T_{\max}$ .
- 20           50. The method of claim 47 wherein the improved pharmacokinetics comprises an  
increase in  $C_{\max}$  of the substance compared to subcutaneous injection.
51. The method of claim 47 wherein the improved pharmacokinetics is a decrease in  
 $T_{\text{lag}}$ .
- 25           52. The method of claim 47 wherein the improved pharmacokinetics is an enhanced  
absorption rate.
53. The method of Claim 46 wherein the device has a fluid driving means including  
a syringe, infusion pump, piezoelectric pump, electromotive pump, electromagnetic pump,  
30   or Belleville spring.
53. The method of Claim 46 wherein the dermal access means comprises one or  
more hollow microcannula having a length of from about 0.5 to about 1.7 mm- mm.
- 35           54. The method of Claim 46 wherein said dermal access means comprises one or  
more hollow microcannula having an outlet with an exposed height between 0 and 1 mm.

WO 02/02179

PCT/US01/20782

- 5            55. A method for delivering a bioactive substance to a subject comprising:  
contacting the skin of a subject with a device having a dermal-access means for accurately  
targeting the dermal space of the subject      with an efficacious amount of the bioactive  
substance at a rate of 1nL/min. to 200 mL/min.
- 10           56. The method of claim 55 wherein the rapid onset pharmacokinetics of the  
bioactive substance is substantially improved relative to subcutaneous injection.
57. The method of claim 56 wherein the bioavailability is increased.
- 15           58. The method of claim 56 wherein the pharmacokinetics is a decreased  $T_{max}$ .
59. The method of claim 56 wherein the pharmacokinetics is an increased  $C_{max}$ .
60. The method of claim 56 wherein the pharmacokinetics is a decreased  $T_{lag}$ .
- 20           61. The method of claim 56 wherein the pharmacokinetics is an enhanced absorption  
rate.
62.     The method of Claim 55 wherein the dermal access means has one or more  
hollow microcannula that inserts into the skin of said subject to a depth of from about 0.5 to  
25     about-2.0 mm.
63. The method of Claim 55 wherein the dermal access means has one or more  
hollow microcannula having an outlet with an exposed height between 0 and 1 mm.

1/7



WO 02/02179

PCT/US01/20782

2/7





FIG. 2



WO 02/02179

PCT/US01/20782

3/7



WO 02/02179

PCT/US01/20782

4/7




FIG. 4

5/7




FIG. 5

WO 02/02179

PCT/US01/20782

6/7




FIG. 6

SUBSTITUTE SHEET (RULE 26)

7/7




FIG. 7

## INTERNATIONAL SEARCH REPORT

International Application No  
PC1/US 01/20782

**A. CLASSIFICATION OF SUBJECT MATTER**  
IPC 7 A61M37/00

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)  
IPC 7 A61M A61B

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the International search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 3 814 097 A (GANDERTON D ET AL) 4 June 1974 (1974-06-04) abstract; claim 1	38-45
A	WO 99 64580 A (GEORGIA TECH RES INST) 16 December 1999 (1999-12-16) abstract; claim 7	38-45
A	US 5 879 326 A (ANDERSON R ROX ET AL) 9 March 1999 (1999-03-09) abstract	38-45
A	US 2 619 962 A (ROY ROSENTHAL SOL) 2 December 1952 (1952-12-02) column 2, line 31,32	38-45

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

\* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the International filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the International filing date but later than the priority date claimed

- \*T\* later document published after the International filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- \*A\* document member of the same patent family

Date of the actual completion of the International search

1 November 2001

Date of mailing of the International search report

12/11/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Nielsen, M

## INTERNATIONAL SEARCH REPORT

International Application No  
PCT/US 01/20782

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 3814097	A	04-06-1974	GB 1408925 A	08-10-1975
			AR 195415 A1	08-10-1973
			AU 5145073 A	25-07-1974
			BE 795384 A1	13-08-1973
			CH 560544 A5	15-04-1975
			DE 2305989 A1	23-08-1973
			DK 131270 B	23-06-1975
			FR 2172168 A1	28-09-1973
			IT 995037 B	10-11-1975
			JP 48093192 A	03-12-1973
			LU 67007 A1	19-04-1973
			NL 7301843 A	16-08-1973
			ZA 7300459 A	31-10-1973
WO 9964580	A	16-12-1999	AU 4561699 A	30-12-1999
			EP 1086214 A1	28-03-2001
			WO 9964580 A1	16-12-1999
US 5879326	A	09-03-1999	AU 5740496 A	11-12-1996
			WO 9637155 A1	28-11-1996
US 2619962	A	02-12-1952	NONE	